

We claim:

1. A liposome composition comprising DOTAP and at least one cholesterol or cholesterol derivative.

2. The liposome composition according to claim 1, further comprising a biologically-active agent, thereby forming a sandwich liposome.

3. The sandwich liposome composition according to claim 2 wherein the composition has a  $\rho$  value equal to 2.

4. The liposome composition according to claim 2, wherein the biologically-active agent is a nucleic acid.

5. The liposome composition according to claim 4 further comprising, adding a targeting ligand thereby decorating exterior surface of said sandwich liposome with the ligand.

6. A DNA-sandwich liposome composition comprising a structure having lipid bilayers and DNA molecules positioned between two or more sandwich liposomes, wherein  $\rho = 2$  and a size of 200 - 450 nm.

7. A DNA-sandwich liposome comprising DNA, DOTAP and at least one of a cholesterol or cholesterol derivative.

8. The DNA-sandwich liposome of claim 7 further comprising one or more targeting ligands.

9. A liposome produced by the steps comprising:

i) heating DOTAP and at least one cholesterol or cholesterol derivative forming heated lipid

components;

- ii) sonicating said heated lipid components; and
- iii) extruding lipid components sequentially through filters of decreasing pore size.

10. The liposome of claim 9 further comprising a sandwich liposome, produced by adding a biologically-active agent to the liposomes.

11. The liposome of claim 10 wherein the biologically active agent is DNA, thereby forming a DNA sandwich liposome.

12. The liposome according to claim 11 further comprising, adding a targeting ligand thereby decorating the exterior surface of said DNA-sandwich liposome with the ligand.

13. The liposome according to claim 11 further comprising a second biologically active agent.

14. The liposome of claim 11 wherein the DNA, DOTAP and at least one cholesterol or cholesterol derivative carry a  $\rho$  value of 2.

15. A method for preparing invaginated liposomes comprising the steps of:

- i) heating a mixture of DOTAP and at least one of cholesterol or cholesterol derivative forming heated lipid component;
- ii) sonicating said heated lipid components; and
- iii) extruding lipid components sequentially through filters of decreasing pore size forming invaginated liposomes.

16. The method of claim 15, further comprising adding DNA to said invaginated liposomes forming DNA-sandwich liposomes.